



SFRP4 gene

secreted frizzled related protein 4

Normal Function

The *SFRP4* gene provides instructions for making a protein called secreted frizzled-related protein 4 (SFRP4). This protein blocks (inhibits) a process called Wnt signaling. Wnt signaling plays an important role in the development of several tissues and organs throughout the body. In particular, regulation of this signaling process by SFRP4 is critical for normal bone development and remodeling. Bone remodeling is a normal process in which old bone is broken down and new bone is created to replace it. The SFRP4 protein also plays a role in the development of fatty (adipose) tissue.

Health Conditions Related to Genetic Changes

Dupuytren contracture


Pyle disease

At least four *SFRP4* gene mutations have been found in individuals with a bone disorder called Pyle disease. This condition is characterized by a bone abnormality in which the ends (metaphyses) of the long bones in the arms and legs are abnormally wide, resembling a boat oar or paddle. Other bones may also be abnormal in Pyle disease, including the collar bones (clavicles), ribs, and bones in the fingers and hands. The *SFRP4* gene mutations are thought to lead to production of an abnormally short SFRP4 protein with impaired function, or they result in no SFRP4 protein production at all. Studies suggest that loss of functional SFRP4 dysregulates Wnt signaling, which disrupts normal bone development and remodeling. Abnormal bone formation leads to the characteristics of Pyle disease.

Chromosomal Location

Cytogenetic Location: 7p14.1, which is the short (p) arm of chromosome 7 at position 14.1

Molecular Location: base pairs 37,905,932 to 37,916,923 on chromosome 7 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- frizzled protein, human endometrium
- FRP-4
- FRPHE
- PYL
- secreted frizzled-related protein 4 precursor
- secreted frizzled-related protein 4; secreted frizzled-related protein 4
- sFRP-4

Additional Information & Resources

Educational Resources

- Endotext (2000): Bone Remodeling and its Relationship to Bone Quantity
https://www.ncbi.nlm.nih.gov/books/NBK279134/#_epidem-patho-osteopo_toc-bone-remodeling-and-its-relationship-to-bone-quantity_
- Madame Curie Bioscience Database (2000): Secreted Frizzled-Related Protein (sFRPs)
https://www.ncbi.nlm.nih.gov/books/NBK6536/#_A16295_

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28SFRP4%5BTIAB%5D%29+OR+%28secreted+frizzled+related+protein+4%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5BIa%5D+AND+human%5Bmh%5D+AND+%22last+1440+days%22%5Bdp%5D>

OMIM

- SECRETED FRIZZLED-RELATED PROTEIN 4
<http://omim.org/entry/606570>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/SFRP4ID42277ch7p14.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=SFRP4%5Bgene%5D>
- HGNC Gene Family: Secreted frizzled-related proteins
<http://www.genenames.org/cgi-bin/genefamilies/set/733>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=10778
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/6424>
- UniProt
<http://www.uniprot.org/uniprot/Q6FHJ7>

Sources for This Summary

- Galada C, Shah H, Shukla A, Girisha KM. A novel sequence variant in SFRP4 causing Pyle disease. J Hum Genet. 2017 Jan 19. doi: 10.1038/jhg.2016.166. [Epub ahead of print]
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/28100910>
- Haraguchi R, Kitazawa R, Mori K, Tachibana R, Kiyonari H, Imai Y, Abe T, Kitazawa S. sFRP4-dependent Wnt signal modulation is critical for bone remodeling during postnatal development and age-related bone loss. Sci Rep. 2016 Apr 27;6:25198. doi: 10.1038/srep25198.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27117872>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4846872/>
- OMIM: SECRETED FRIZZLED-RELATED PROTEIN 4
<http://omim.org/entry/606570>

- Simsek Kiper PO, Saito H, Gori F, Unger S, Hesse E, Yamana K, Kiviranta R, Solban N, Liu J, Brommage R, Boduroglu K, Bonafé L, Campos-Xavier B, Dikoglu E, Eastell R, Gossiel F, Harshman K, Nishimura G, Girisha KM, Stevenson BJ, Takita H, Rivolta C, Superti-Furga A, Baron R. Cortical-Bone Fragility--Insights from sFRP4 Deficiency in Pyle's Disease. *N Engl J Med*. 2016 Jun 30; 374(26):2553-62. doi: 10.1056/NEJMoa1509342.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/27355534>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5070790/>
- Visweswaran M, Schiefer L, Arfuso F, Dilley RJ, Newsholme P, Dharmarajan A. Wnt antagonist secreted frizzled-related protein 4 upregulates adipogenic differentiation in human adipose tissue-derived mesenchymal stem cells. *PLoS One*. 2015 Feb 25;10(2):e0118005. doi: 10.1371/journal.pone.0118005.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/25714610>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4340908/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/SFRP4>

Reviewed: March 2017

Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services